蝶と蛾 Trans. lepid. Soc. Japan 52 (3): 168-170, June 2001

Summer disappearance of territoriality in the satyrine butterfly Lethe sicelis

Jun-Ya IDE

Department of Zoology, Graduate School of Science, Kyoto University, Kitashirakawa Oiwake-cho, Sakyo-ku, Kyoto, 606-8502 Japan; e-mail: ide@terra.zool.kyoto-u.ac.jp

Abstract Lethe sicelis displays territorial behaviour related to mating. Territorial behaviour was observed in June and September, but not in July and August, although L. sicelis was observed. It is suggested that high air temperatures prevent L. sicelis from holding territory in summer.

Key words Air temperature, *Lethe sicelis*, seasonal change, territorial behaviour.

Introduction

The behaviours of butterflies often change seasonally with changing air temperature (Hidaka, 1975; Ide, 2000a), since butterflies are ectotherms. The males of many butterfly species hold territories for mating (Shreeve, 1992) and territorial behaviour also changes seasonally, both in time (Bitzer & Shaw, 1995; Fric & Konvička, 2000) and place (Shreeve, 1987). The satyrine butterfly *Lethe diana* (Butler) shows dramatic changes in mate-locating behaviour with season; males of *L. diana* display territorial behaviour in May–June and September-October, but not in July–August (Ide, in press).

Lethe sicelis (Hewitson), a congener with L. diana, also displays territorial behaviour (Fukuda et al., 1984). However, no seasonal change in the territoriality of L. sicelis has been reported to date. In this study, I recorded the territorial behaviour of L. sicelis observed for three years, and here describe seasonal changes in its territoriality.


Materials and methods

In the *L. sicelis* flight season from 1998 to 2000, transect censuses were carried out on almost every sunny day (66, 85, and 97 days in 1998, 1999, 2000, respectively). A fixed route of approximately 3 km was established in the Kamigamo Experimental Forest Station, Graduate School of Agriculture, Kyoto University, in the north of Kyoto City, Japan (135°46′E, 35°04′N; 140 m in altitude); this was the same route as reported in Ide (2000*b*). I walked along the route and counted *L. sicelis* seen within an estimated distance of 5 m ahead, above, or on either side of me, in accordance with the method of Ishii (1993). When a butterfly was found holding a territory, it was recorded.

Results and discussion

L. sicelis was observed at the station from June to September every year. Territorial behaviour was observed, mostly around sunlit branch tips that faced open spaces, in June and September, but it was scarcely observed in July and August (Fig. 1). The mean air temperatures on the days when territorial behaviour was observed never exceeded 25.5°C. One territorial male was observed in August, but it was on a cloudy day and the mean air

Summer Disappearance of Territoriality in Lethe sicelis






Fig. 1. Seasonal change in territoriality in *Lethe sicelis*. Territorial behaviour was observed in June and September, but was not observed in July or August, although *L. sicelis* was observed.

Jun-Ya Ide

temperature was low.

The disappearance of territorial behaviour in summer has also been reported in *L. diana* (Ide, in press). A male *L. diana* displays territorial behaviour only when his body temperature is maintained within a suitable range. In summer, the thermal environment of the territory is too hot, and *L. diana* cannot display territorial behaviour (Ide, unpublished). *L. sicelis* also did not display territorial behaviour when it was hot, which suggests that high temperatures prevent *L. sicelis* from displaying territorial behaviour. The mate-locating behaviour of male *L. sicelis* in July and August, when they do not hold territories, is not known. Further research is necessary to clarify the mate-locating behaviour of *L. sicelis* during all seasons.

Acknowledgements

I thank the Kamigamo Experimental Forest Station, Graduate School of Agriculture, Kyoto University, for permission to conduct this research.

References

- Bitzer, R. J. & K. C. Shaw, 1995. Territorial behaviour of the red admiral, *Vanessa atalanta* (Lepidoptera: Nymphalidae) I. The role of climatic factors and early interaction frequency on territorial start time. *J. Insect Behav.* 8: 47-66.
- Fric, Z. & M. Konvička, 2000. Adult population structure and behaviour of two seasonal generations of the Europian Map Butterfly, *Araschnia levana*, species with seasonal polyphenism (Nymphalidae). *Nota lepid.* 23: 2-25.
- Fukuda, H., Hama, E., Kuzuya, T., Takahashi, A., Takahashi, M., Tanaka, B., Tanaka, H., Wakabayashi, M. & Y. Watanabe, 1984. *The Life Histories of Butterflies in Japan* 4. xxii, 65-373 pp., 64 pls. Hoikusha, Osaka. (In Japanese).
- Hidaka, T., 1975. Why does the Butterfly Fly. 178 pp. Iwanami-shoten, Tokyo. (In Japanese).
- Ide, J.-Y., 2000a. Seasonal change in flight behaviour of the satyrine butterfly *Lethe diana* (Lepidoptera: Nymphalidae). *Ent. Sci.* 3: 591–596.
- ______, 2000b. Seasonal change in factors affecting spatial microdistribution in a population of the satyrine butterfly Lethe diana. Trans. lepid. Soc. Japan 52: 13-24.
- ——— (in press). Mating behaviour and light conditions cause seasonal changes in the dispersal pattern of the satyrine butterfly *Lethe diana*. *Ecol. Ent.*
- Ishii, M., 1993. Transect counts of butterflies. *In* Yata, O. and K. Ueda (Eds), Decline and Conservation of Butterflies in Japan II. *Yadoriga* (spec. Issue): 91-101 (in Japanese with English summary).
- Shreeve, T. G., 1987. The mate location behaviour of the male speckled wood butterfly, *Pararge aegeria*, and the effect of phenotypic differences in hind-wing spotting. *Anim. Behav.* **35**: 682-690.
- , 1992. Adult behaviour. *In* Dennis, R. L. H. (Ed.), *The Ecology of Butterflies in Britain*: 22-45. Oxford University Press, Oxford.

摘 要

ヒカゲチョウのなわばり行動の季節変化 (井出純哉)

ヒカゲチョウ (ジャノメチョウ亜科)の雄は配偶行動としてなわばり行動を行う. 3 年間の調査の結果、本種の成虫は6 月から9 月まで出現し6 月と9 月にはなわばり行動が見られるが、7-8 月には見られないことがわかった. 同属のクロヒカゲと同様、暑い時期には体温が上がり過ぎないように、なわばり行動をやめている可能性がある.

(Accepted March 26, 2001)

Published by the Lepidopterological Society of Japan, 5-20, Motoyokoyama 2, Hachioji, Tokyo, 192-0063 Japan